

Day : Monday
Date: 7/7/2003Time:
19:26:30PALM INTRANET**Inventor Name Search Result**

Your Search was:

Last Name = ZHANG

First Name = JIANPING

Application#	Patent#	Status	Date Filed	Title	Inventor Name
<u>60458818</u>	Not Issued	020	03/28/2003	PROCESS AND APPARATUS FOR CONTROLLING FLOW IN A MULTIPHASE REACTOR	ZHANG, JIANPING
<u>60344229</u>	Not Issued	020	12/28/2001	WATER REMOVAL IN FISCHER-TROPSCH PROCESSES	ZHANG, JIANPING
<u>60344228</u>	Not Issued	020	12/28/2001	METHOD FOR REDUCING WATER CONCENTRATION IN A MULTI-PHASE COLUMN REACTOR	ZHANG, JIANPING
<u>60307742</u>	Not Issued	020	07/25/2001	MINIMIZING THE VOLUME OR MAXIMIZING THE PRODUCTION RATE OF SLURRY BUBBLE REACTORS BY USING LARGE GAS FLOW RATES AND MODERATE SINGLE PASS CONVERSION	ZHANG, JIANPING
<u>10402498</u>	Not Issued	019	03/28/2003	GAS AGITATED MULTIPHASE CATALYTIC REACTOR WITH REDUCED BACKMIXING	ZHANG, JIANPING
<u>10395912</u>	Not Issued	019	03/24/2003	COMMERCIAL FISCHER-TROPSCH REACTOR	ZHANG, JIANPING
<u>10320311</u>	Not Issued	030	12/16/2002	WATER REMOVAL IN FISCHER-TROPSCH PROCESSES	ZHANG, JIANPING
<u>10315571</u>	Not Issued	030	12/10/2002	METHOD FOR REDUCING WATER CONCENTRATION IN A MULTI-PHASE	ZHANG, JIANPING

				COLUMN REACTOR	
<u>10238008</u>	Not Issued	030	09/09/2002	GAS AGITATED MULTIPHASE REACTOR WITH STATIONARY CATALYST SOLID PHASE	ZHANG, JIANPING
<u>10205215</u>	Not Issued	030	07/25/2002	MINIMIZING THE VOLUME OR MAXIMIZING THE PRODUCTION RATE OF SLURRY BUBBLE REACTORS BY USING LARGE GAS FLOW RATES AND MODERATE SINGLE PASS CONVERSION	ZHANG, JIANPING
<u>10193357</u>	Not Issued	030	07/11/2002	METHOD FOR REDUCING THE MAXIMUM WATER CONCENTRATION IN A MULTI-PHASE COLUMN REACTOR	ZHANG, JIANPING
<u>10023258</u>	Not Issued	030	12/14/2001	SLURRY BED REACTOR	ZHANG, JIANPING
<u>09559936</u>	Not Issued	094	04/27/2000	DEPOLARIZED SEMICONDUCTOR LASER SOURCES	ZHANG, JIANPING

Inventor Search Completed: No Records to Display.

Search Another:
Inventor

Last Name

Zhang

First Name

Jianping

Search

To go back use Back button on your browser toolbar.

Back to [PALM](#) | [ASSIGNMENT](#) | [OASIS](#) | [Home page](#)

Welcome to STN International! Enter x:x

LOGINID:sssptal202jxp

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	Jun 03	New e-mail delivery for search results now available
NEWS	4	Aug 08	PHARMAMarketLetter(PHARMAML) - new on STN
NEWS	5	Aug 19	Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN
NEWS	6	Aug 26	Sequence searching in REGISTRY enhanced
NEWS	7	Sep 03	JAPIO has been reloaded and enhanced
NEWS	8	Sep 16	Experimental properties added to the REGISTRY file
NEWS	9	Sep 16	CA Section Thesaurus available in CAPLUS and CA
NEWS	10	Oct 01	CASREACT Enriched with Reactions from 1907 to 1985
NEWS	11	Oct 24	BEILSTEIN adds new search fields
NEWS	12	Oct 24	Nutraceuticals International (NUTRACEUT) now available on STN
NEWS	13	Nov 18	DKILIT has been renamed APOLLIT
NEWS	14	Nov 25	More calculated properties added to REGISTRY
NEWS	15	Dec 04	CSA files on STN
NEWS	16	Dec 17	PCTFULL now covers WP/PCT Applications from 1978 to date
NEWS	17	Dec 17	TOXCENTER enhanced with additional content
NEWS	18	Dec 17	Adis Clinical Trials Insight now available on STN
NEWS	19	Jan 29	Simultaneous left and right truncation added to COMPENDEX, ENERGY, INSPEC
NEWS	20	Feb 13	CANCERLIT is no longer being updated
NEWS	21	Feb 24	METADEx enhancements
NEWS	22	Feb 24	PCTGEN now available on STN
NEWS	23	Feb 24	TEMA now available on STN
NEWS	24	Feb 26	NTIS now allows simultaneous left and right truncation
NEWS	25	Feb 26	PCTFULL now contains images
NEWS	26	Mar 04	SDI PACKAGE for monthly delivery of multifile SDI results
NEWS	27	Mar 20	EVENTLINE will be removed from STN
NEWS	28	Mar 24	PATDPAFULL now available on STN
NEWS	29	Mar 24	Additional information for trade-named substances without structures available in REGISTRY
NEWS	30	Apr 11	Display formats in DGENE enhanced
NEWS	31	Apr 14	MEDLINE Reload
NEWS	32	Apr 17	Polymer searching in REGISTRY enhanced
NEWS	33	Jun 13	Indexing from 1947 to 1956 added to records in CA/CAPLUS
NEWS	34	Apr 21	New current-awareness alert (SDI) frequency in WPIDS/WPINDEX/WPIX
NEWS	35	Apr 28	RDISCLOSURE now available on STN
NEWS	36	May 05	Pharmacokinetic information and systematic chemical names added to PHAR
NEWS	37	May 15	MEDLINE file segment of TOXCENTER reloaded
NEWS	38	May 15	Supporter information for ENCOMPPAT and ENCOMPLIT updated
NEWS	39	May 16	CHEMREACT will be removed from STN
NEWS	40	May 19	Simultaneous left and right truncation added to WSCA
NEWS	41	May 19	RAPRA enhanced with new search field, simultaneous left and right truncation
NEWS	42	Jun 06	Simultaneous left and right truncation added to CBNB
NEWS	43	Jun 06	PASCAL enhanced with additional data

NEWS 44 Jun 20 2003 edition of the FSTA Thesaurus is now available
NEWS 45 Jun 25 HSDB has been reloaded

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT
MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),
AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 13:49:45 ON 04 JUL 2003

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 13:49:57 ON 04 JUL 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 4 Jul 2003 VOL 139 ISS 2

FILE LAST UPDATED: 3 Jul 2003 (20030703/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s au 200137521/pn

L1 0 AU 200137521/PN
(AU2001037521/PN)

=> s wo 200172928/pn

L2 1 WO 200172928/PN
(WO2001072928/PN)

=> d 12 iall

L2 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2003 ACS
 ACCESSION NUMBER: 2001:730909 CAPLUS
 DOCUMENT NUMBER: 135:275198
 TITLE: Method for producing hydrocarbons from syngas in a three-phase reactor
 INVENTOR(S): Schweitzer, Jean-Marc; Galtier, Pierre; Hugues, Francois; Maretto, Cristina
 PATENT ASSIGNEE(S): Institut Francais du Petrole, Fr.; Eni S.p.A.; Agip Petroli S.p.A.
 SOURCE: PCT Int. Appl., 21 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 INT. PATENT CLASSIF.:
 MAIN: C10G002-00
 CLASSIFICATION: 51-11 (Fossil Fuels, Derivatives, and Related Products)
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001072928	A1	20011004	WO 2001-FR595	20010228 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
FR 2806736	A1	20010928	FR 2000-3925	20000327
FR 2806736	B1	20020510		
EP 1268709	A1	20030102	EP 2001-909933	20010228
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
NO 2002004608	A	20020926	NO 2002-4608	20020926
US 2003109590	A1	20030612	US 2002-239395	20020926
PRIORITY APPLN. INFO.:			FR 2000-3925	A 20000327
			WO 2001-FR595	W 20010228

ABSTRACT:

The invention concerns a method for the synthesis of hydrocarbons by reacting a mixt. comprising at least carbon monoxide and hydrogen in the presence of a catalyst carried out in a three-phase reactor, wherein the liq. Peclet no. ranges between 0 (excluded) and .apprx.10, at a gas surface speed Ug preferably <35 cm./s, to promote the transfer of the gas into the liq. phase and avoid too much attrition of catalyst grains.

SUPPL. TERM: fuel gas manufg syngas catalyst
 INDEX TERM: Fuel gas manufacturing
 Synthesis gas
 (method for producing hydrocarbons from syngas in a three-phase reactor)
 INDEX TERM: Rare earth oxides
 ROLE: CAT (Catalyst use); USES (Uses)
 (method for producing hydrocarbons from syngas in a three-phase reactor)
 INDEX TERM: 1314-23-4, Zirconia, uses 1344-28-1, Alumina, uses
 7440-48-4, Cobalt, uses 7631-86-9, Silica, uses
 13463-67-7, Titania, uses

ROLE: CAT (Catalyst use); USES (Uses)
(method for producing hydrocarbons from syngas in a
three-phase reactor)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD.

REFERENCE(S):
(1) Exxon Research Engineering Co; EP 0450861 B 1991 CAPLUS
(2) Iglesia, E; US 5348982 A 1994
(3) Sasol Chemicals Europ Ltd; WO 9903574 A 1999 CAPLUS



Generate Collection

Print

L4: Entry 24 of 55

File: USPT

May 26, 1992

DOCUMENT-IDENTIFIER: US 5116879 A

** See image for Certificate of Correction **

TITLE: Process using a supported catalyst for hydrocarbon synthesis

Brief Summary Text (5):

The reaction to convert carbon monoxide and hydrogen mixtures (so called syngas) to higher hydrocarbons over metallic catalysts has been known since the turn of the century. This reaction is commonly referred to as the Fischer-Tropsch or F-T synthesis. The F-T synthesis was exploited commercially during WWII in Germany. By 1944 a total of nine F-T plants were operating in Germany, primarily using a catalyst composed of cobalt, magnesium oxide, thorium oxide, and kieselguhr, in the relative composition 100:5:8:200. Later, most of the thorium was replaced by magnesia, primarily for economic reasons. Currently, commercial Fischer-Tropsch plants, which use a precipitated iron-based catalyst which contains various promoters to improve the stability and product distribution, are operating in South Africa.

Brief Summary Text (6):

The common F-T catalysts are nickel, cobalt, and iron. Nickel was probably the first substance to be recognized as capable of catalyzing the reaction of syngas to hydrocarbons, producing mainly methane (see, for example, R. B. Anderson, The Fischer-Tropsch Synthesis, Academic Press (1984), p. 2). Iron and cobalt are able to produce higher chain hydrocarbons and are, thus, preferred as catalysts for the production of liquid hydrocarbons. However, other metals are also capable of catalyzing the conversion of synthesis gas. Among the Group VIII metals, ruthenium is a very active catalyst for the formation of hydrocarbons from syngas. Its activity at low temperatures is higher than that of Fe, Co, or Ni, and it produces a large amount of heavy hydrocarbons. At high pressures, it produces a large amount of high molecular weight waxes. Other metals which are highly active, such as rhodium, yield high amounts of oxygenated materials (see Ichikawa, Chemtech, 6, 74 (1982)). Osmium has been found to be moderately active, while Pt, Pd, and Ir exhibit low activity (see Pichler, Advances in Catalysis, vol. IV, Academic Press, N.Y. (1952), R. B. Anderson, The Fischer-Tropsch Synthesis, supra and Vannice, Journal of Catalysis, 50, 228-236).

Brief Summary Text (7):

Other metals that have been investigated include rhenium, molybdenum, and chromium, but these exhibit very low activity with most of the product being methane (see R. B. Anderson, The Fischer-Tropsch Synthesis, supra).

Brief Summary Text (9):

Combinations of cobalt with non-F-T active metals have also been reported for the conversion of synthesis gas to specific products and, in some cases, at specific conditions. In Nakaoji, U.S. Pat. No. 3,988,344, the combination of cobalt with a second Group VIII metal and tungsten is claimed for the enhanced production of methane from synthesis gas. Knifton, U.S. Pat. No. 4,390,734 and Japanese Kokai 57/130932 describe the combination of Co and Rh for the production of oxygenated products, such as glycols or aldehydes. Fischer-Tropsch catalysts consisting of combinations of cobalt with either platinum or palladium, supported on a variety of solids, including alumina, have been reported by Sapienza et al., U.S. Pat. No. 4,396,539. These catalysts, however, relied on preparation from the metal carbonyls in order to form solid solutions on the surface of the solid support and were distinguished by an x-ray impermeable layer covering the support, thereby resulting in a catalyst exhibiting a unique x-ray diffraction pattern in which the structure of the solid support was completely masked by the metallic components. In

and more preferably from 1000 to 10,000 cm.sup.3 /g/h, where gaseous hourly space velocity is defined as the volume of gas (measured at standard temperature and pressure) fed per unit weight of catalyst per hour.

Detailed Description Text (25):

The hydrocarbon products from Fischer-Tropsch synthesis are distributed from methane to high boiling compounds according to the so called Schulz-Flory distribution, well known to those skilled in the art. The Schulz-Flory distribution is expressed mathematically by the Schulz-Flory equation:

Detailed Description Text (28):

It is well known, and also shown in the following examples, that the metals from the group consisting of platinum, iridium, and rhodium are alone low activity catalysts for Fischer-Tropsch synthesis producing products which are predominantly methane, or in the case of rhodium, oxygenates.

CLAIMS:

1. A process for the production of hydrocarbons comprising the step of contacting a synthesis gas feed comprised of hydrogen and carbon monoxide with a catalyst consisting essentially of an amount of cobalt catalytically active in a Fischer-Tropsch synthesis, and at least one second metal selected from the group consisting of iridium, rhodium, and mixtures thereof composited on an alumina support, said second metal being present in relatively lesser amounts than the cobalt content of the catalyst, at a temperature, gas hourly space velocity and a pressure useful for promoting a Fischer-Tropsch Synthesis.

3. A process for the production of hydrocarbons comprising the step of contacting a synthesis gas feed comprised of hydrogen and carbon monoxide with a catalyst consisting essentially of an amount of cobalt catalytically active in a Fischer-Tropsch synthesis, at least one second metal selected from the group consisting of iridium, rhodium, and mixtures thereof, and a promoter, wherein said promoter is an oxide of a metal selected from the group consisting of periodic groups IIIB, IVB, and VB, the lanthanides, the actinides, magnesium, and mixtures thereof, composited on an alumina support, said second metal being present in relatively lesser amounts than the cobalt content of the catalyst, said promoter ranging from about 0.1 to about 5 wt % of the catalyst, at a temperature, gas hourly space velocity and a pressure useful for promoting a Fischer-Tropsch Synthesis.

particular, when the catalysts of the Sapienza et al. patent are supported on alumina, they are distinguished by the complete absence of any x-ray diffraction peaks in the 2 .theta. range of 65 to 70 degrees. In x-ray diffraction .theta. equals the angle of refraction.

Brief Summary Text (11):

In a series of European patent applications (EP 110449, EP 142888, EP 167215, and EP 188304), Shell International described an improved F-T catalyst comprised of cobalt promoted by at least one of the metals in the group consisting of zirconium, titanium, and chromium, preferably supported on silica, alumina, or silica-alumina. The addition of Group VIII noble metals to zirconia-promoted cobalt catalysts was claimed in European patent application EP 221598, which teaches improved activity upon addition of platinum to a cobalt catalyst already promoted with zirconia. Thus, the Shell work shows that the addition of a Group VIII metal to a Fischer-Tropsch catalyst is only useful when the catalyst already incorporates a well known promoter, such as zirconia, as a main component of the catalyst.

Brief Summary Text (14):

It has been found in accordance with the present invention that synthesis gas comprising hydrogen and carbon monoxide can be converted to liquid hydrocarbons by using a process which includes the step of contacting such a synthesis gas feed over a catalyst comprising amounts of cobalt catalytically active in a Fischer-Tropsch Synthesis and at least one loading-insensitive second metal selected from the group consisting of platinum, iridium, and rhodium composited on an alumina support. The finished catalyst exhibits a positive x-ray diffraction pattern having peaks in the 2.theta. range of about 65 to 70 degrees, where .theta. is the angle of refraction. The second metal is present in relatively lesser amounts than the cobalt content. The catalyst preferably contains from about 5 to 60 wt % cobalt and has a content of the second metal between about 0.1 and 50 wt % of the cobalt content of the catalyst. The alumina preferably is gamma alumina.

Brief Summary Text (15):

It has been found that the addition of one or more of the metals from the group consisting of platinum, iridium, and rhodium to a catalyst consisting predominantly of cobalt supported on alumina results in greatly enhanced activity for the conversion of synthesis gas to hydrocarbons. Platinum, iridium, and rhodium are not very active as Fischer-Tropsch catalysts on their own, and it has surprisingly been found that the addition of the other non-Fischer-Tropsch noble metals to a cobalt catalyst does not result in greatly improved activity. The improvement upon addition of platinum, iridium, or rhodium to a cobalt catalyst is surprisingly not observed when the catalytic components are distended on supports other than alumina, for example silica or titania.

Detailed Description Text (2):

The catalyst used in the process of the present invention comprises as the active catalytic ingredients cobalt and one or more metals from the group consisting of platinum, iridium, and rhodium supported on alumina with the second metal present in a relatively smaller amount than cobalt. The finished catalyst exhibits a positive x-ray diffraction pattern having peaks in the 2 .theta. range of about 65 to 70 degrees. This catalyst has been found to be highly active for the conversion of synthesis gas, a mixture of hydrogen and carbon monoxide, into a mixture of predominately paraffinic hydrocarbons. As indicated above, it has long been known that cobalt is an active catalyst for the F-T synthesis. It is also known that the addition of ruthenium to a cobalt catalyst gives improved activity, but ruthenium is known to be an active Fischer-Tropsch metal on its own. In our invention, it has been found that among the non-Fischer-Tropsch metals in Group VIII of the Periodic Table, some of these metals produce enhanced activity when added to a supported cobalt catalyst without the need to form solid solutions on the surface of the support, while others of these metals give no improvement. Although a number of supports have been studied in the present work and the improvement of this invention has only been observed with alumina, the discovery of another support exhibiting a similar effect would not be entirely surprising.

Detailed Description Text (20):

The reaction temperature is suitably between 150.degree. and 300.degree. C.; and more preferably between 175.degree. and 250.degree. C. The total pressure can be from atmospheric to around 100 atmospheres, preferably between 1 and 30 atmospheres. The gaseous hourly space velocity, based on the total amount of synthesis gas feed, is preferably between 100 and 20,000 cm.sup.3 of gas per gram of catalyst per hour;